



FEED THE FUTURE

The U.S. Government's Global Hunger & Food Security Initiative

Feed the Future Innovation Lab for Livestock Systems

Burkina Faso: Animal-Source Food Production and Marketing

February 2017

The Management Entity at the University of Florida



USAID
FROM THE AMERICAN PEOPLE



UF | IFAS
UNIVERSITY of FLORIDA

Acknowledgement

The Animal-Source Food Production and Marketing brief was prepared by Andrew Noss PhD under the supervision of Geoffrey Dahl, PhD, Leader of the Area of Inquiry on Animal-Source Food Production and Marketing.

This brief is a work in progress. It will be updated with additional information collected in the future.

This brief is made possible with the generous support of the people of the United States (US) through the US Agency for International Development (USAID) under the Feed the Future Initiative. The contents in this brief are the responsibility of the University of Florida and do not necessarily reflect the views of USAID or the United States government, and its partners in Feed the Future.

1. Introduction

Estimated annual growth rates for Burkina Faso livestock numbers have been 2% (goats, sheep, and chickens) and 3% (cattle, pigs) from 2005 to 2014. In 2014, the number of goats exceeded 13.8 million head, followed by sheep at 9.3 million, cattle at over 9.1 million, and pigs at 2.3 million. The national camel herd was only about 18,000 animals (MRA, 2015). Of these animals, women raise goats more than men do, while principally men raise the others (INERA, 2015). Chickens are by far the most common domestic animal, 33.7 million, though guinea fowl are also numerous at 8.4 million. Women manage 70% of poultry of which more than half (60%) is also marketed by women, principally in local markets (MRA, 2007a). Egg production doubled from 2012-2014, to over 120,000, though only 10% of eggs are sold (MRA, 2015; CRS, 2014b; INERA, 2015). The Sahel region has the largest number of cattle, goats, sheep, and camels; while Centre-Ouest region has the largest number of pigs, chickens and guinea fowl (Figure 1).

Table 1. Livestock numbers by region in 2014, thousands (MRA, 2015).

Region	Cattle	Goats	Sheep	Pigs	Chickens	Guinea fowl
Boucle du Mouhoun	803	1195	761	271	4163	1115
Cascades	676	226	236	53	884	324
Centre	156	338	224	186	1330	196
Centre-Est	427	1073	731	218	2793	521
Centre-Nord	512	1280	1027	76	2436	296
Centre-Ouest	706	1684	1032	457	4898	2120
Centre-Sud	318	756	400	125	2534	871
Est	1034	1470	950	130	2535	445
Hauts-Bassins	1509	813	850	265	4226	896
Nord	415	1246	921	136	2868	588
Plateau Central	324	861	577	113	2071	280
Sahel	1868	2329	1301	4	1383	225
Sud-Ouest	343	620	268	312	1632	591

Based on research in the country, the Institute for Agricultural and Environmental Research (INERA) and the Food and Agricultural Organization of the United Nations (FAO) and other institutions have published extensively on ways to improve the productivity of livestock in Burkina Faso. This brief summarizes information on production systems, forages and feeds, markets, processing, and genetic improvement.

2. Production Systems

The Ministry of Livestock Resources (MRA, 2007b; 2011a; 2011b) categorizes livestock production systems as traditional extensive or commercial, with numerous sub-categories, most of which combine multiple livestock species.

Traditional extensive production systems

Transhumant production systems

Transhumant production systems (9% of livestock producers, 16% of cattle and 6% of goats and sheep) are operated mainly by Peulh pastoralists, within Burkina Faso but also extending north to and from Mali and Niger, and extending south to the coastal countries. Average cattle herd size is 60 animals and the Sudanese Peulh zebu breed is favored. In Boucle du Mouhoun and East Sahel, transhumant herds of sheep and goats (Peulh, Gourmantché, Bella races) are also found. Transhumance is either long-distance, > 100 km, or short distance. Livestock in this system depends on natural pastures, and there is no integration

with crop production, except that herds use fields post-harvest to feed on the crop residues. Livestock receive little or no inputs of shelter, minerals, or feed supplements. Only lactating animals may be supplemented in the dry season. Forage is rarely produced or stored. Livestock also depend greatly on surface water and on wells, and face difficulties in the dry season. Other challenges for transhumance systems are the conversion of natural pasture lands and transhumance corridors to crops, lack of secure land and resource tenure, and conflicts with farmers (Kagoné, 2001; MRA, 2007b; 2011a; 2011b; Diallo & Boundaogo, 2011; FAO, 2012; Traore, 2012).

Sedentary extensive systems (within one day's walk)

Today these are the predominant production systems in Burkina Faso, practiced by 85% of livestock producers, and involving 79% of cattle, 91% of goats and sheep, 92% of pigs and camels, and 98% of chickens and guinea fowl (MRA, 2011a; 2011b; Guissou et al., 2013).

- a. Agro-pastoral sedentary systems are used to convert crop surplus to livestock. These are widely practiced by most ethnic groups and by white-collar workers. One version involves raising sedentary cattle away from crops, using natural pastures, while the other integrated system keeps animals with crops (traction cattle, donkeys, horses as well as sheep and goats). In these systems livestock benefit from dry-season feeding of crop residues and brans (MRA, 2007b; Diallo & Boundaogo, 2011). Poultry are kept in mixed flocks of up to 15 birds, with no to minimal surveillance, little management of reproduction, only rudimentary shelter, and no vaccination, though some water and supplementary feed are provided (MRA, 2007a).
- b. Urban and peri-urban sedentary systems are practiced by residents of Ouagadougou and Bobo-Dioulasso who raise goats, sheep, cows, and chickens. Animals are pastured year-round in open city spaces, as well as on fallows and rangelands at the cities' edges. Owners also produce forage for livestock, and provide supplemental feed such as brewery wastes or cottonseed cakes for ruminants, and cereal grains and brans for poultry. Manure is provided as compensation (not sold) to owners of crop fields where animals forage, and sometimes is used in family gardens. Most animals are vaccinated and de-parasitized, but reproduction is not managed. Herd size averages 8 goats and sheep (Diallo & Boundaogo, 2011; Kaboré et al., 2011; Amadou et al., 2012; FAO, 2012; Dossa et al., 2015; Gnanda et al., 2016). Poultry flocks can be up to 50 birds and production is for income. Sturdy and clean shelters are provided, as are water and feed (MRA, 2007a).

Commercial semi-intensive or intensive production systems

Cattle fattening and milk production

- a. Family cattle fattening operations handle up to three animals at a time, providing several types of feed and mineral supplements: agro-industrial byproducts (cottonseed or other), grain straw, natural forages, salt. Principally men undertake cattle fattening, and animals are sold in local markets (MRA, 2007b; Diallo & Boundaogo, 2011; Traore, 2012; CRS, 2014a; 2014b).
- b. Commercial fattening and milk production systems around Ouagadougou and Bobo-Dioulasso are operated by civil servants, traders or private companies. They use native cattle breeds (Peulh, Goudali and Azawak) as well as cross-breeding of these or with exotics. Operators produce their own forages, cut and dry natural forages, and purchase crop residues and agro-industrial byproducts as well as mineral supplements. They provide good veterinary care and utilize artificial insemination techniques. Fattening operations handle 10-100 animals at a time (Hamadou & Sanon, 2005; Diallo & Boundaogo, 2011; Sanon et al., 2014).

Sheep fattening

Goat fattening is less common because market demand is low (CRS, 2014a).

- a. Family sheep fattening is principally done by women—using Bali Bali, Djallonké or mixed-breed sheep—up to five animals at a time (MRA, 2007b). Animals are sold principally in local markets. Sheep receive a variety of feeds including cowpea and peanut greens, grain straw, natural forages, cottonseed, local bran (MRA, 2007b; Diallo & Boundaogo, 2011; Traore, 2012; CRS, 2014a; 2014b).
- b. Commercial sheep fattening is undertaken by traders, and more recently by women’s groups sponsored by NGOs or government projects (MRA, 2007b).

Other semi-intensive and intensive production systems

Other semi-intensive and intensive production systems include all pigs in Burkina Faso, and are increasing for poultry and milk production to meet growing demand. These are located near urban areas and are operated by civil servants and entrepreneurs (MRA, 2011a).

3. Milk Production

Milk production was traditionally the responsibility of women (and especially the Peulh, Mossi, Tuareg and Maure pastoralist peoples), and their principal income source, but with livestock further from residences and with government and project interventions, men have taken over milk marketing and income (Hamadou & Sanon, 2005). Although productivity is very low (0.5-2.0 liters per cow), dairy products contribute on average 32% of farm income, and are important in each of the livestock management systems described above: transhumant, sedentary extensive, and semi-intensive to intensive (FAO, 2012). Numbers of milk transformation units (UTL) have fluctuated between 100 and 200 from 2007-2014, with the highest numbers in Centre and Hauts Bassins Regions, where urban demand is highest from Ouagadougou and Bobo-Dioulasso, respectively; as well as in Sahel Region where cattle, goats and sheep are most abundant. Milk production has likewise fluctuated over the same period, exceeding 3.5 million liters of processed milk in 2014, with nearly half the production in Hauts-Bassins Region alone (MRA, 2015). The Islamic Development Bank is financing from 2014-2018 the implementation of the Société de Production de Lait (SOPROLAIT), with an expected production capacity of 30,000 liters of UHD milk per year (Traore, 2012).

4. Feeds and Forages

Overall, insufficient and poor quality forage and feed greatly limit productivity. The livestock production systems described above depend very heavily on natural pastures (fresh grasses, and some woody vegetation in the dry season) for ruminants and on local products (household waste but also cereal grains and brans) for pigs and poultry (Table 2, Appendices 1-3). Cultivated forages are virtually absent, though FAO has been promoting them since at least 1950 (Kagoné, 2001; Guissou et al., 2013). Markets in urban areas do supply forage products, generally natural forages or crop residues, at least on a seasonal basis (Kaboré et al., 2012; Tamini et al., 2014). However, commercialization is limited by the lack of transport and storage facilities, difficulties with drying natural forage, bush fires, and the distance to harvest sites (Tamini et al., 2014). Forage collection methods, often taking the entire plant, can denude the landscape, while feed quality is poor under rudimentary processing (silage or urea treatment is rare) and storage methods that rarely enclose or shade the fodder (CRS, 2014a; ILRI & CGIAR, 2015). Natural forages are also declining as pastoral lands are converted to crops (INERA, 2015), yet crop residues (rice, millet, sorghum, and maize straws, peanut, Bambara groundnut, and cowpea foliage) remain underutilized (Kagoné, 2001; MRA, 2007b; Traore, 2012; Sanon & Traore, 2014; Sanon et al., 2014; Gnanda et al., 2016). In order to increase forage production, INERA is producing seed for forage crops, which MRA

provides to livestock producers at a subsidized rate (80% of actual cost), together with forage harvesting equipment (ILRI & CGIAR, 2015).

More than 50 small private producers now supply agro-industrial byproducts, mainly cottonseed cakes and cereal brans, while several large private or public-private companies. Companies supplying raw materials and/or producing feeds include MINOFA, SN CITEC, SOFAB, ICO, SOBA, SOFITEX, GMB, SN SOSUCO, Unité ASUDEDEC, LAPHAVET, CPAVI, Genol BF, and SOFIB Huilerie. Cotton grain, molasses, and brewery waste are also utilized (Sanon et al., 2014). However, their high cost limits them to urban and peri-urban production systems and intensive livestock operations such as the fattening systems (Diallo & Boundaogo, 2011; Amadou et al., 2012; Traore, 2012; CRS, 2014a; 2014b; Sanon & Traore, 2014; Sanon et al., 2014; INERA, 2015). MRA is making them available at a subsidized rate (30% of actual cost) (ILRI & CGIAR, 2015). Only SOFAB (Société de Fabrique d'Aliment pour Bétail) markets a formula specific to goats and sheep (Tamini et al., 2014). SOFAB alone has a production capacity of 10,000 tons of feed per year but only operates at half capacity (A. Ouedraogo, personal communication).

Table 2. Principal food sources by livestock species (%) (MRA, 2011a).

	Natural pasture	Agro-industrial byproducts	Crop residues	Cultivated forages	Hay	Local products
Cattle	83	2	12	<1	<1	2
Sheep	80	2	15	<1	<1	3
Goats	84	1	12	<1	<1	2
Pigs	9	2	4	<1	3	81
Camels	51	49	<1	0	0	<1
Chickens	0	<1	6	<1	0	93
Guinea fowl	0	<1	5	0	<1	95

5. Exports and Marketing

In terms of ASF, Burkina Faso officially exports about 10 times the value of what it imports, and over 90% of imports are milk products (MRA 2011b). Burkina Faso is a major exporter of livestock, particularly to its coastal neighbors to the south. Côte d'Ivoire is the most important destination for sheep and chickens, while Ghana is for goats, and Nigeria is for cattle. Important numbers of ruminants are also exported to Niger, the majority of which are likely re-exported from there. Some meat is exported—peaking at 11 million CFA francs in 2010 (approximately \$20,000) but declining to less than 2 million in 2012 (\$3600). Ninety-five percent of meat exports, 142,000 kg in 2012, are to Côte d'Ivoire and to Gabon (MRA, 2015). But most exports are live animals: official export figures are 500,000 sheep, 400,000 goats, 340,000 cattle, and 300,000 chickens; with cattle representing 80% of the 5.2 billion CFA in 2011 earnings. Official export numbers likely underestimate the exports. For example, Jossierand (2013) estimated that export statistics may underestimate outflows by a factor of 4 to 8.

Three categories of internal livestock markets operate in Burkina Faso (MRA, 2007b; Traore, 2012; Guissou et al., 2013; Tamini et al., 2014):

- Production or primary collection markets in the rural areas where livestock is raised and where herders sell 1-2 animals at a time, with no livestock handling infrastructure (water sources, corrals, scales). Intermediary buyers build up herds for transfer to other markets, moving the animals on foot or by truck.
- Redistribution markets are medium-sized markets in towns on transit corridors; some are traditional with no infrastructure for handling livestock, some are modern with corrals and truck loading chutes.

- Final or export markets. Fada N’Gourma, the largest, is a big modern market. Others include Ouagadougou and Bobo-Dioulasso, among others.

Although the typology of the markets is important and largely applies, the distinctions are not clear cut as e.g., a trader may stop to purchase animals from a primary collection market, in search for cheaper prices or to fill up a truck.

6. Processing

The MRA recognizes nearly 400 abattoirs and slabs around the country, where licensed inspectors certify animal handling procedures for 1,200,000 goats, 400,000 sheep, 250,000 cattle, and 180,000 pigs slaughtered annually in these legal facilities (MRA 2015). Only four public abattoirs have cold-storage capabilities (Ouagadougou, Bobo-Dioulasso, Dédougou, and Ouahigouya), and one private one (Tamini et al., 2014), though these are not up to international standards. Many other animals of these species, and all poultry, are slaughtered at home or in non-certified abattoirs, representing fully two-thirds of meat consumed in Burkina Faso (MRA, 2007a; Traore 2012; Fadiga & Fall, 2014; Tamini et al., 2014; ILRI & CGIAR, 2015).

In turn, virtually all meat is consumed fresh, sold directly to consumers, to meat-grillers, and to restaurants (MRA, 2007b; Tamini et al., 2014). Only two modern meat-processing plants operate, in Ouagadougou and Bobo-Dioulasso respectively (ILRI & CGIAR, 2015).

7. Genetic Improvement

Although they are resistant to local diseases and environmental conditions, native breeds are low in productivity (MRA, 2007b). In northern Burkina Faso, the larger Peulh or Sahel sheep and goat breeds predominate, and in the south and center of the country the smaller Djallonké and Mossi breeds or cross-breeds (Tamini et al., 2014). The four principal local breeds of chickens are Peulh, Dori, Kondé, and Central Burkina Faso gray (MRA, 2007a).

Various initiatives for genetic improvement have been undertaken, but have generally been limited. The positive features of local breeds, including disease resistance and adaptation to local environmental conditions, could also be lost or diluted by exotic breeds (Tamini et al., 2014). The International Atomic Energy Agency is financing the project BKF5014 “Improving the Productivity of Small Ruminants through Diet, Health and Identification of Genetic Markers for Selection and Breeding Management” from 2014-2017. The objective of the project is “to contribute to improving the productivity and profitability of small ruminant farms in Burkina Faso by applying genetic characterization and artificial insemination for breeding and utilizing local feed resources to improve nutrition and medicinal plants to control parasites” (<https://www.iaea.org/projects/tc/bkf5014>). The government of Luxembourg is also financing the project BKF/017, “Support for the dissemination of the Azawak Zebu”, to improve the productivity of this local cattle breed (<https://luxdev.lu/en/activities/project/BKF/017>).

Literature Cited

- Amadou, H., Dossa, L.H., Lompo, D.J.P., Abdulkadir, A. & Schlecht. 2012. A comparison between urban livestock production strategies in Burkina Faso, Mali and Nigeria in West Africa. *Tropical Animal Health Production* 44:1631–1642.
- Catholic Relief Services. 2014a. Adapting to the Sahel’s changing climate: local approaches. An economic and technical feasibility analysis of adaptation techniques in Niger, Burkina Faso and Mali. Baltimore, MD, CRS.
- . 2014b. Building livelihoods on the frontlines of climate change: Identifying market opportunities and agricultural value chains in the Sahel regions of Niger, Burkina Faso and Mali. Baltimore, MD, CRS.
- Diallo, L., and Boundaogo, M. 2011. Les principales filières agro-sylvo-pastorales de la province du Sanmatenga. Ouagadougou, Burkina Faso: SNV.
- Dossa, L.H., Sangaré, M., Buerkert, A. & Schlecht, E. 2015. Production objectives and breeding practices of urban goat and sheep keepers in West Africa: regional analysis and implications for the development of supportive breeding programs. *SpringerPlus* 4:281. DOI 10.1186/s40064-015-1075-7
- Fadiga, M. & Fall, A. 2014. A scoping study of small ruminants value chains in the ECOWAS countries. Ouagadougou, Burkina Faso, ILRI
- FAO, ECOWAS. 2012. The cross-border transhumance in West Africa Proposal for Action Plan. Rome, FAO.
- Gnanda, B.I., Wereme N’Diaye, A., Sanon, H.O., Somda, J. & Nianogo, J.A. 2016. Rôle et place de la chèvre dans les ménages du Sahel burkinabé. *Tropicicultura* 34(1):10-25.
- Guissou, R., Ilboudo, F., Cissé, K., and Honoré, Z., 2013. Analyse des incitations et pénalisations pour les bovins au Burkina Faso. Technical Note Series. Rome : SPAAA (Suivi des Politiques Agricoles et Alimentaires en Afrique Project), FAO.
- Hamadou, S. & Sanon, Y. 2005. Synthèse bibliographique sur les filières laitières au Burkina-Faso. Document de travail No. 3. Réseau de Recherche et d’Echanges sur les Politiques Laitière. www.repol.sn
- ILRI & CGIAR. 2015. Revue des innovations technologiques et des options politiques relatives aux chaînes de valeur des petits ruminants au Burkina Faso. 1ere version provisoire. Rapport Synthèse. Ouagadougou, Burkina Faso.
- INERA (Institut de l’Environnement et de Recherches Agricoles), ILRI, CGIAR. 2015. Evaluation des chaînes de valeur des Petits Ruminants dans les sites de Thiou (Ouahigouya, Yatenga), Bani (Dori, Séno), Bama, (Bobo Dioulasso, Houet) et Kaya, Sanematenga), au Burkina Faso. 1ere version provisoire. Ouagadougou, Burkina Faso.
- Josserand, H. 2013. Assessment of volumes and value of regionally traded staple commodities. Paper prepared for the Food Across Borders Conference, Accra, 29-31 January 2013. http://www.inter-reseaux.org/IMG/pdf/Josserand_-_Assessment_of_ATP_E-ATP_Trade_Data.pdf.

- Kaboré, A., Traoré, A., Gnanda, B.I., Nignan, M., Tamboura, H.H., & Belem, A.M.G. 2011. Constraints of small ruminant production among farming systems in periurban area of Ouagadougou, Burkina Faso (West Africa). *Advances in Applied Science Research*, 2(6):588-594.
- Kaboré, A., Traoré, A., Gnanda, B.I., Ouédraogo, P., Tamboura, H.H., & Belem A.M.G. 2012. Diversity and characteristics of green natural fodders sold in the city of Ouagadougou in Burkina Faso, West Africa. *International Research Journal of Agricultural Science and Soil Science* 2(8): 377-385.
- Kagoné, H. 2001. Profil fourrager. Ouagadougou, Burkina Faso: MRA, FAO.
- Ministry of Livestock Resources (MRA). 2007a. Diganostic de la sous-filière de l'aviculture traditionnelle au Burkina Faso. Ouagadougou, Burkina Faso.
- . 2007b. Plans d'actions pour le développement des filières bétail-viande et petits ruminants. Première partie : Diagnostic des filières – Rapport final. Ouagadougou, Burkina Faso.
- . 2011a. Contribution de l'élevage à l'économie et a la lutte contre la pauvreté, les déterminants de son développement. Ouagadougou, Burkina Faso.
- . 2011b. Document de plaidoyer du sous-secteur de l'élevage. Ouagadougou, Burkina Faso.
- . 2015. Annuaire des statistiques de l'élevage 2013-2014. Ouagadougou, Burkina Faso.
- Sanon, H.O., & Traore, L. 2014. Typologie d'ateliers d'embouche ovine au Burkina Faso (Ouest) et performances des unités pilotes. *Livestock Research for Rural Development* 26(10):1-9.
- Sanon, H.O., Drabo, A., Sangare, Kiendrebeogo, T., & Gomgnibou, A. 2014. Caractérisation des pratiques d'embouche bovine dans l'Ouest du Burkina Faso. *International Journal of Biological and Chemical Sciences* 8(2):536-550.
- Tamini, L.D., Fadiga, M.L., & Sorgho, Z. 2014. Chaines de valeur des petits ruminants au Burkina Faso: Analyse de situation. Addis Ababa, Ethiopia: ILRI (International Livestock Research Institute).
- Traore, F. 2012. Republique du Burkina Faso: Rapport de pré diagnostic des filières Bétail – Viande et sésame. Ouagadougou, Burkina Faso: African Agribusiness and Agro-industries Development Initiative, IFAD, FAO, UNIDO.

Appendix 1: Forage species by agro-ecological zone (Kagoné, 2001)

Agro-ecological zones (annual rainfall)	Grasses: Native and Cultivated	Legumes: Native and Cultivated
Sahel (< 400 mm)	<i>Andropogon gayanus</i> <i>Cenchrus ciliaris</i> cv. <i>Biloela</i> <i>Cenchrus ciliaris</i> cv. <i>Gayndah</i> <i>Pennisetum pedicellatum</i>	<i>Alysicarpus ovalifolius</i> <i>Macroptilium atropurpureum</i> <i>Macroptilium lathyroides</i> Cultivated: <i>Stylosanthes hamata</i> Forage cowpea
Sub-Sahel (400–600 mm)	<i>Andropogon gayanus</i> <i>Pennisetum pedicellatum</i> Cultivated: Forage sorghum Forage maize	<i>Alysicarpus ovalifolius</i> <i>Lablab niger</i> <i>Macroptilium atropurpureum</i> <i>Mucuna</i> sp. Cultivated: Forage cowpea
North-Sudan (600–900 mm)	<i>Andropogon gayanus</i> <i>Panicum maximum</i> <i>Pennisetum pedicellatum</i> <i>Rottboellia exaltata</i> Cultivated: Forage sorghum	<i>Aeschynomene histrix</i> <i>Crotalaria intermedia</i> <i>Crotalaria juncea</i> <i>Lablab niger</i> <i>Mucuna</i> sp. <i>Mucuna pruriens</i> cv <i>utilis</i> Cultivated: <i>Stylosanthes gracilis</i> <i>Stylosanthes guianensis</i> Forage cowpea
South-Sudan (> 900 mm)	<i>Andropogon gayanus</i> <i>Brachiaria ruziziensis</i> <i>Cenchrus ciliaris</i> cv. <i>Biloela</i> <i>Chloris gayana</i> <i>Echinochloa stagnina</i> <i>Melinis minutiflora</i> <i>Panicum maximum</i> <i>Pennisetum purpureum</i> cv <i>Napier</i> Cultivated: Forage sorghum Forage maize	<i>Aeschynomene histrix</i> <i>Cajanus cajan</i> <i>Centrosema</i> spp. <i>Dol</i> <i>Lablab niger</i> <i>Neonotonia wightii</i> <i>Macroptilium atropurpureum</i> <i>Macroptilium lathyroides</i> <i>Mucuna</i> sp. <i>Phaseolus aureus</i> Cultivated: <i>Stylosanthes hamata</i> cv <i>Verano</i> <i>Stylosanthes guianensis</i>

Appendix 2: Other Native Forages (Kagoné, 2001; ILRI & CGIAR, 2015)

Grasses	Legumes	Other
<i>Acroceras amplexans</i>	<i>Acacia ehenbergiana, A. laeta, A.</i>	<i>Adansonia digitata</i>
<i>Andropogon pseudacpricus</i>	<i>nilotica, A. raddiana, A. senegal, A.</i>	<i>Aerva javanica</i>
<i>Anogeissus leiocarpus</i>	<i>seyal</i>	<i>Balanites aegyptiaca</i>
<i>Aristida mutabilis, A. hordaceae, A.</i>	<i>Azelia africana</i>	<i>Boerhavia erecta</i>
<i>kerstingii, A. adscensionis</i>	<i>Bauhinia rufescens</i>	<i>Butyrospermum</i>
<i>Burkea africana</i>	<i>Cassia obtusifolia</i>	<i>paradoxum</i>
<i>Butyrospermum paradoxum</i>	<i>Dolichos lablab</i>	<i>Combretum glutinosum</i>
<i>Cenchrus biflorus, C. prieurii</i>	<i>Indigofera aspera</i>	<i>Evolvulus alsinoides</i>
<i>Cymbopogon schoenanthus</i>	<i>Isoberlinia doka</i>	<i>Gardenia tenuifolia</i>
<i>Dactyloctenium aegyptium</i>	<i>Molluga nudicaulis</i>	<i>Kyaha senegalensis</i>
<i>Digitaria horizontalis</i>	<i>Parkia biglobosa</i>	<i>Limeum viscosum, L.</i>
<i>Diheteropogon hagerupii</i>	<i>Piliostigma reticulatum</i>	<i>pterocarpum</i>
<i>Echinochloa pyramidalis</i>	<i>Pterocarpus erinaceus, P. lucens</i>	<i>Terminalia avicennioides,</i>
<i>Eragrostis tremuula</i>	<i>Tamarindus indica</i>	<i>T. macroptera</i>
<i>Hyparrhenia cyanescens</i>	<i>Zornia glochidiata</i>	<i>Tribulus terrestris</i>
<i>Imperata cylindrica</i>		
<i>Loudetia togoensis</i>		
<i>Loudetiopsis scaettae</i>		
<i>Mitragyna inermis</i>		
<i>Oryza barthii</i>		
<i>Panicum anabaptistum, P. laetum</i>		
<i>Schiizachyrium exile, S. sanguineum, S.</i>		
<i>brevifolium</i>		
<i>Schoenefeldia gracilis</i>		
<i>Vetiveria nigriflora</i>		

Appendix 3: Principal feed sources for small ruminants (INERA, 2015)

Natural pasture	Cultivated forage	Purchased feed
Grasses		
<i>Andropogon gayanus</i>	Cowpea greens	Cottonseed cakes
<i>Eragrostis tremula</i>	Peanut greens	Cotton seeds
<i>Loudetia togoensis</i>	Sorghum / millet / maize stalks	Corn bran
<i>Pennisetum pedicellatum</i>	Hibiscus stalks	Sorghum stalks
<i>Shoenfeletia gracilis</i>	Sweet potato leaves	Native grass (dry)
Legumes		
<i>Acacia nilotica, A. albida, A. seyal,</i>	Mucuna	Cubed bran
<i>A. machrostachyn, A. ataxaeanta</i>	Rice straw	Rice straw
<i>Alisicarpus ovalifolices</i>		
<i>Afzelia africana</i>		
<i>Bauhinia rufescens</i>		
<i>Cassia sieberiana</i>		
<i>Eichrostachys glomerata</i>		
<i>Zornia glachidiata</i>		
Other		
<i>Adansonia digitata</i>		
<i>Anogeisus leiocarpus</i>		
<i>Combretum aculeatum</i>		
<i>Dalbergia sissoo</i>		
<i>Grewia bicolor, G. molle</i>		
<i>Ipomoea oocarpa</i>		
<i>Kyaha senegalensis</i>		
<i>Lannea microcarpa, L. acida</i>		
<i>Securinega virosa</i>		
<i>Ziziphus mauritiana</i>		